

Next Nearest-Neighbor Correlation Functions of the Spin-1/2 XXZ Chain at Massive Region

Minoru Takahashi,^{1,*} Go Kato,^{2,†} and Masahiro Shiroishi^{1,‡}

¹*Institute for Solid State Physics, University of Tokyo, Kashiwa, Chiba 277-8571, Japan*

²*Department of Physics, Graduate School of Science,
University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan*

The second neighbor correlation functions of the spin- $\frac{1}{2}$ XXZ chain in the ground state are expressed in the form of three dimensional integrals. We show that these integrals can be reduced to one-dimensional ones and thereby evaluate the values of the next nearest-neighbor correlation functions for $\Delta > 1$.

PACS numbers: 75.10.Jm, 75.50.Ee, 02.30.Ik

I. INTRODUCTION

The spin-1/2 XXZ chain is one of the fundamental models in the study of the low-dimensional magnetism. The Hamiltonian is given by

$$H = \sum_{j=1}^N \{S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + \Delta S_j^z S_{j+1}^z\}, \quad (1)$$

where $S = \sigma/2$ and σ are Pauli matrices. The model can be solved by Bethe ansatz method and diverse physical properties have been investigated with varying anisotropy parameter Δ [1, 2, 3]. For $\Delta = 1$ we know the second neighbor correlation [4]. Jimbo et al [5, 6, 7] obtained the multiple integral representation for the arbitrary correlation functions of the XXZ chain for $1 \geq \Delta > -1$ and $\Delta > 1$. Later Kitanine, Maillet and Terras[8] generalized to magnetized case using the Bethe wave function. But this multiple integral is very com-

*mtaka@issp.u-tokyo.ac.jp

†kato@monet.phys.s.u-tokyo.ac.jp

plicated and they only reproduce the nearest neighbor correlations. Recently Boos et al succeeded to calculate these multiple integrals for emptiness formation probability (EFP)

$$P(m) = \langle \prod_{j=1}^m (S_j^z + \frac{1}{2}) \rangle, \quad (2)$$

at $\Delta = 1, m = 3, 4, 5, 6$ [9, 10, 11, 12]. The case $m = 3$ reproduces the result of the second neighbor correlation. The same method is applied to the third neighbor correlations at $\Delta = 1$ [13]. In the previous paper we calculated the second neighbor correlation functions for massless case $-1 < \Delta \leq 1$ and integral is simplified to the one dimensional integral[14]. In this paper we give the second neighbor correlation in massive region $1 < \Delta$. The correlation formula contains the elliptic functions. But we can show that the final result also contain only one-dimensional integral.

‡siroisi@issp.u-tokyo.ac.jp

II. MULTIPLE INTEGRAL FORMULA

A. Massless case

The correlation function in massless case is given by,

$$F(\epsilon_1, \epsilon_2, \dots, \epsilon_m) = \frac{\langle \psi_g | E_{\epsilon_1}^{\epsilon_1} E_{\epsilon_2}^{\epsilon_2} \dots E_{\epsilon_m}^{\epsilon_m} | \psi_g \rangle}{\langle \psi_g | \psi_g \rangle} = (-1)^s \left(-\frac{\pi}{\zeta}\right)^{\frac{m(m+1)}{2}} \prod_{j=1}^{s'} \int_{-\infty-i\zeta}^{\infty-i\zeta} \frac{d\lambda_j}{2\pi} \prod_{j=s'+1}^m \int_{-\infty}^{\infty} \frac{d\lambda_j}{2\pi} \prod_{a>b} \frac{\sinh \frac{\pi}{\zeta}(\lambda_a - \lambda_b)}{\sinh(\lambda_a - \lambda_b - i\zeta)} \\ \times \prod_{j \in \alpha^+} \frac{\sinh^{j-1}(\mu'_j + 3i\frac{\zeta}{2}) \sinh^{m-j}(\mu'_j + i\frac{\zeta}{2})}{(\cosh \frac{\pi}{\zeta} \mu'_j)^m} \prod_{j \in \alpha^-} \frac{\sinh^{j-1}(\mu_j - i\frac{\zeta}{2}) \sinh^{m-j}(\mu_j + i\frac{\zeta}{2})}{(\cosh \frac{\pi}{\zeta} \mu_j)^m}, \quad (3)$$

where

$$\Delta = \cos \zeta, \quad \{\lambda_1, \dots, \lambda_m\} = \{\mu'_{j'_{max}}, \dots, \mu'_{j'_{min}}, \mu_{j_{min}}, \dots, \mu_{j_{max}}\}, \\ \alpha^+ = \{j; 1 \leq j \leq m, \epsilon_j = +\}, \text{card}(\alpha^+) = s', \quad \text{Max}_{j \in \alpha^+}(j) \equiv j'_{max}, \text{Min}_{j \in \alpha^+}(j) = j'_{min}, \\ \alpha^- = \{j; 1 \leq j \leq m, \epsilon_j = -\}, \text{card}(\alpha^-) = s, \quad \text{Max}_{j \in \alpha^-}(j) \equiv j_{max}, \text{Min}_{j \in \alpha^-}(j) = j_{min}. \quad (4)$$

If we put $\lambda_j \rightarrow \lambda_j - i\zeta$ for $j = 1, \dots, s'$, these equations are written as follows

$$F(\epsilon_1, \epsilon_2, \dots, \epsilon_m) = (-1)^m \left(-\frac{\pi}{\zeta}\right)^{\frac{m(m+1)}{2}} \times \prod_{j=1}^m \int_{-\infty}^{\infty} \frac{d\lambda_j}{2\pi} \frac{\prod_{a>b} \sinh \frac{\pi}{\zeta} (\lambda_a - \lambda_b)}{\prod_{j=1}^m \cosh^m \frac{\pi \lambda_j}{\zeta}} \frac{\prod_{j=1}^m \sinh^{y_j-1} (\lambda_j + i\frac{\zeta}{2}) \sinh^{m-y_j} (\lambda_j - i\frac{\zeta}{2})}{\prod_{a>b} \sinh (\lambda_a - \lambda_b - i f_{ab} \zeta)}, \quad (5)$$

where

$$f_{ab} = (1 + \text{sign}[(s' + 1/2 - a)(s' + 1/2 - b)])/2, \\ y_1 > y_2, \dots > y_{s'}, \quad \epsilon_{y_i} = +, \quad y_{s'+1} > y_{s'+2} > \dots > y_m, \quad \epsilon'_{m+1-y_i} = -. \quad (6)$$

If we put $\lambda_j \rightarrow x_j \zeta$ this is written as

$$\prod_{j=1}^m \int_{-\infty}^{\infty} dx_j W_m(x_1, \dots, x_m) \frac{\prod_{j=1}^m \sinh^{y_j-1} \zeta (x_j + \frac{i}{2}) \sinh^{m-y_j} \zeta (x_j - \frac{i}{2})}{\zeta^{m(m-1)/2} \prod_{a>b} \sinh \zeta (x_a - x_b - i f_{ab})}. \quad (7)$$

The weight function of the integral is common for all the correlation functions:

$$W_m(x_1, \dots, x_m) = \frac{\prod_{a>b} (-\pi) \sinh \pi (x_a - x_b)}{\prod_{j=1}^m 2 \cosh^m \pi x_j} = D_m \left(\prod_{j=1}^m \frac{1}{2} \text{sech}(\pi x_j) \right) = D_m \left(\prod_{j=1}^m W_1(x_j) \right), \quad (8)$$

where D_m is antisymmetric differential operator defined by

$$D_m = \frac{1}{\prod_{r=1}^{m-1} r!} \prod_{j>k} \left(\frac{\partial}{\partial x_j} - \frac{\partial}{\partial x_k} \right). \quad (9)$$

Especially in the limit $\zeta \rightarrow 0$ or $\Delta = 1$ we have

$$\prod_{j=1}^m \int_{-\infty}^{\infty} dx_j W_m(x_1, \dots, x_m) \frac{\prod_{j=1}^m (x_j + \frac{i}{2})^{y_j-1} (x_j - \frac{i}{2})^{m-y_j}}{\prod_{a>b} (x_a - x_b - i f_{ab})}. \quad (10)$$

These integrals are done analytically up to $m = 4$ for $\Delta = 1$ [13], and up to $m = 3$ for $-1 < \Delta < 1$ [14].

B. Massive case

For massive case integral formula for correlation function contains elliptic functions [5, 7, 8]

$$F(\epsilon_1, \epsilon_2, \dots, \epsilon_m) = (i)^{s-s'} C_m \prod_{j=1}^{s'} \int_{-\pi/2-i\phi}^{\pi/2-i\phi} \frac{d\lambda_j}{2\pi} \prod_{j=s'+1}^m \int_{-\pi/2}^{\pi/2} \frac{d\lambda_j}{2\pi} \vartheta_2 \left(\sum_{j=1}^m (\lambda_j + i\phi/2) \right) \prod_{a>b} \frac{\vartheta_1(\lambda_a - \lambda_b)}{\sin(\lambda_a - \lambda_b - i\phi)} \\ \times \prod_{j \in \alpha^+} \frac{\sin^{j-1}(\mu'_j + 3i\frac{\phi}{2}) \sin^{m-j}(\mu'_j + i\frac{\phi}{2})}{\vartheta_1^m(\mu'_j + i\phi/2)} \prod_{j \in \alpha^-} \frac{\sin^{j-1}(\mu_j - i\frac{\phi}{2}) \sin^{m-j}(\mu_j + i\frac{\phi}{2})}{\vartheta_1^m(\mu_j + i\phi/2)}, \quad (11)$$

where

$$\Delta = \cosh \phi, \quad q = e^{-\phi}, \quad C_m = \prod_{n=1}^{\infty} \left(\frac{1-q^{2n}}{1+q^{2n}} \right)^2 \left[2q^{1/4} \prod_{n=1}^{\infty} (1-q^{2n})^3 \right]^{\frac{m(m+1)}{2}-1} = \frac{(\vartheta'_1(0))^{\frac{m(m+1)}{2}}}{\vartheta_2(0)},$$

$$\vartheta_1(x) = q^{1/4} \sum_{n=0}^{\infty} (-1)^n q^{n(n+1)/2} \sin(2n+1)x, \quad \vartheta_2(x) = q^{1/4} \sum_{n=0}^{\infty} q^{n(n+1)/2} \cos(2n+1)x. \quad (12)$$

Then these equations are written as follows

$$F(\epsilon_1, \epsilon_2, \dots, \epsilon_m) = \prod_{j=1}^m \int_{-\pi/2}^{\pi/2} \frac{d\lambda_j}{2\pi} U_m(\lambda_1, \dots, \lambda_m) \frac{\prod_{j=1}^m \sin^{y_j-1}(\lambda_j + i\frac{\phi}{2}) \sin^{m-y_j}(\lambda_j - i\frac{\phi}{2})}{\prod_{a>b} \sin(\lambda_a - \lambda_b - if_{ab}\phi)}, \quad (13)$$

$$U_m(\lambda_1, \dots, \lambda_m) = i^m C_m \frac{\prod_{a>b} \vartheta_1(\lambda_a - \lambda_b)}{\prod_{j=1}^m \vartheta_1^m(\lambda_j + i\phi/2)} \vartheta_2(\sum_{j=1}^m (\lambda_j + i\phi/2)), \quad (14)$$

where f_{ab} and y_j are given by (6). In Appendix A we show that

$$U_m(\lambda_1, \dots, \lambda_m) = \frac{1}{\prod_{r=1}^{m-1} r!} \prod_{j>k} \left(\frac{\partial}{\partial \lambda_j} - \frac{\partial}{\partial \lambda_k} \right) U_1(\lambda_1) U_1(\lambda_2) \dots U_1(\lambda_m). \quad (15)$$

If we put $\lambda_j \rightarrow x_j \phi$ this is written as

$$\prod_{j=1}^m \int_{-Q/2}^{Q/2} dx_j \mathbf{W}_m(x_1, \dots, x_m) \frac{\prod_{j=1}^m \sin^{y_j-1} \phi(x_j + i\frac{1}{2}) \sin^{m-y_j} \phi(x_j - i\frac{1}{2})}{\phi^{m(m-1)/2} \prod_{a>b} \sin \phi(x_a - x_b - if_{ab})}, \quad (16)$$

$$Q = \pi/\phi, \quad \mathbf{W}_m(x_1, \dots, x_m) = \phi^{m(m-1)/2} U_m(\phi x_1, \dots, \phi x_m). \quad (17)$$

We investigate the properties of $\mathbf{W}_1$,

$$\mathbf{W}_1(x) = i \frac{\vartheta'_1(0) \vartheta_2(\phi(x + i/2))}{\vartheta_2(0) \vartheta_1(\phi(x + i/2))} = \frac{K_l}{\pi Q} \text{dn}\left(\frac{2K_l x}{Q}, l\right), \quad (18)$$

where $\text{dn}(z, l)$ is Jacobian elliptic function with

$$\frac{K(\sqrt{1-l^2})}{K(l)} = \frac{\phi}{\pi}, \quad K(l) = \int_0^{\pi/2} \frac{1}{\sqrt{1-l^2 \sin^2 k}}.$$

This function has poles at $x = kQ + i(l - \frac{1}{2})$ with residue $(-1)^l i$ and zeros at $\lambda = (k - \frac{1}{2})Q + i(l + \frac{1}{2})$. On the other hand $\sum_{n=-\infty}^{\infty} \frac{1}{2} \text{sech}(\pi(x + nQ))$ has the same poles and zeroes. Then we have

$$\mathbf{W}_1(x) = \sum_{n=-\infty}^{\infty} W_1(x + nQ). \quad (19)$$

Using (15) we have

$$\mathbf{W}_m(x_1, \dots, x_m) = D_m \mathbf{W}_1(x_1) \mathbf{W}_1(x_2) \dots \mathbf{W}_1(x_m). \quad (20)$$

Then we have

$$\mathbf{W}_m(x_1, \dots, x_m) = \sum_{n_1, \dots, n_m} W_m(x_1 + n_1 Q, x_2 + n_2 Q, \dots, x_m + n_m Q). \quad (21)$$

The weight function in massless case is trigonometric. But in massive case it becomes elliptic and satisfies the following properties,

$$\mathbf{W}_m(x_1, x_2, \dots, x_j, \dots, x_k, \dots, x_m) = -\mathbf{W}_m(x_1, x_2, \dots, x_k, \dots, x_j, \dots, x_m), \quad (22)$$

$$\mathbf{W}_m(x_1, \dots, x_j, \dots, x_m) = \mathbf{W}_m(x_1, \dots, x_j + Q, \dots, x_m) = -\mathbf{W}_m(x_1, \dots, x_j + i, \dots, x_m), \quad (23)$$

$$\mathbf{W}_m(x_1, \dots, x_j, \dots, x_m) = (-1)^{m(m-1)/2} \mathbf{W}_m(-x_1, \dots, -x_j, \dots, -x_m). \quad (24)$$

We should note that weight function is doubly periodic but the integrand is not doubly periodic. The correlation function has the following properties

$$F(\epsilon_1, \epsilon_2, \dots, \epsilon_m) = F(-\epsilon_1, -\epsilon_2, \dots, -\epsilon_m) = F(\epsilon'_1, \epsilon'_2, \dots, \epsilon'_m) = F(\epsilon_m, \epsilon_{m-1}, \dots, \epsilon_1), \quad (25)$$

$$F(\epsilon_1, \epsilon_2, \dots, \epsilon_{m-1}, +) + F(\epsilon_1, \epsilon_2, \dots, \epsilon_{m-1}, -) = F(\epsilon'_1, \epsilon'_2, \dots, \epsilon'_{m-1}). \quad (26)$$

III. REDUCTION TO ONE DIMENSIONAL INTEGRAL AT $m = 2$ AND 3

At first we consider EFP defined by (2). This is equal to $F(\begin{smallmatrix} +, +, \dots, + \\ +, +, \dots, + \end{smallmatrix})$. As $y_j = m + 1 - j$, $f_{ab} = 1$ and we have the integral representation

$$P(m) = \int_{-Q/2}^{Q/2} d^m x \mathbf{W}_m \frac{\prod_{k=1}^m h^{m-k}(x_k + i/2) h^{k-1}(x_k - i/2)}{\prod_{a>b} h(x_a - x_b - i)}, \quad \text{with } h(x) \equiv \sin(\phi x)/\phi. \quad (27)$$

Recently there have been an increasing number of researches concerning the properties of the EFP. Particularly, in the isotropic limit $\Delta \rightarrow 1$ ($\phi \rightarrow 0$), the general method to evaluate the multiple integral was recently developed by Boos et al[9, 10, 11].

It is important to note that $P(2)$ and $P(3)$ are related to the nearest and next nearest-neighbor correlation functions :

$$\langle S_j^z S_{j+1}^z \rangle = P(2) - 1/4, \quad (28)$$

$$\langle S_j^z S_{j+2}^z \rangle = 2(P(3) - P(2) + 1/8). \quad (29)$$

We also consider transverse correlation functions $\langle S_j^x S_{j+1}^x \rangle$ and $\langle S_j^x S_{j+2}^x \rangle$. Then we should evaluate following four integrals

$$P(2) = \int_{-Q/2}^{Q/2} d^2 x \mathbf{W}_2 \frac{h(x_1 + i/2)h(x_2 - i/2)}{h(x_2 - x_1 - i)}, \quad (30)$$

$$\langle S_j^x S_{j+1}^x \rangle = \frac{1}{2} \int_{-Q/2}^{Q/2} d^2 x \mathbf{W}_2 \frac{h(x_1 + i/2)h(x_2 - i/2)}{h(x_2 - x_1)}, \quad (31)$$

$$P(3) = \int_{-Q/2}^{Q/2} d^3 x \mathbf{W}_3 \frac{h^2(x_1 + \frac{i}{2})h(x_2 + \frac{i}{2})h(x_2 - \frac{i}{2})h^2(x_3 - \frac{i}{2})}{h(x_2 - x_1 - i)h(x_3 - x_1 - i)h(x_3 - x_2 - i)}, \quad (32)$$

$$\langle S_j^x S_{j+2}^x \rangle = \int_{-Q/2}^{Q/2} d^3 x \mathbf{W}_3 \frac{h^2(x_1 + \frac{i}{2})h(x_2 + \frac{i}{2})h(x_2 - \frac{i}{2})h^2(x_3 - \frac{i}{2})}{h(x_2 - x_1)h(x_3 - x_1)h(x_3 - x_2 - i)}. \quad (33)$$

In Appendix C we transform these to the canonical forms. The two-dimensional integrals of (30) and (31) can be reduced to one-dimensional ones,

$$P(2) = \frac{1}{2} + \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x} \frac{1}{2} \left(-\frac{x}{\sin^2 \phi x} + \cot(\phi x) \coth \phi \right), \quad (34)$$

$$\langle S_j^x S_{j+1}^x \rangle = \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x} \frac{1}{4} \left(\frac{x}{\sin^2 \phi x} \cosh \phi - \frac{\cot(\phi x)}{\sinh \phi} \right). \quad (35)$$

These coincide with known results. The main results are that *three-dimensional integrals for $P(3)$ and $\langle S_j^x S_{j+2}^x \rangle$ can also be reduced to one-dimensional integrals*. In other words, we have succeeded in performing the integrals for $P(3)$

TABLE I: Estimated values of correlation function $\langle S_0^z S_2^z \rangle$ from finite rings and our calculation

N	$\Delta = 1.0$	$\Delta = 1.5$	$\Delta = 2.0$	$\Delta = 2.5$	$\Delta = 3.0$	$\Delta = 3.5$	$\Delta = 4.0$
20	0.06135072260	0.1031381904	0.1473943448	0.1795178252	0.1993399825	0.2119381546	0.2204021144
22	0.06123184969	0.1033007174	0.1480705446	0.1799480453	0.1995078997	0.2119982680	0.2204241750
24	0.06114201704	0.1034578190	0.1486278313	0.1802452204	0.1996070522	0.2120292690	0.2204342780
26	0.06107247281	0.1036073511	0.1490859151	0.1804501423	0.1996656632	0.2120452914	0.2204389106
28	0.06101753236	0.1037485416	0.1494616068	0.1805913409	0.1997003229	0.2120535929	0.2204410300
30	0.06097337258	0.1038813030	0.1497692404	0.1806886032	0.1997209063	0.2120579113	0.2204420230
extra.	0.06067894953	0.1061687241	0.1508088848	0.1809161322	0.1999716192	0.2120736609	0.2204679674
exact	0.06067976996	0.1061446753	0.1511283748	0.1809054601	0.1997511670	0.2120626058	0.2204428632

TABLE II: Estimated values of correlation function $\langle S_0^x S_2^x \rangle$ from finite rings and our calculation

N	$\Delta = 1.5$	$\Delta = 2.0$	$\Delta = 2.5$	$\Delta = 3.0$	$\Delta = 3.5$	$\Delta = 4.0$
20	0.04336844424	0.02832886249	0.01874771832	0.01321049421	0.00981186886	0.00757459251
22	0.04318176477	0.02806630512	0.01860734227	0.01315890868	0.00979394498	0.00756813199
24	0.04303078324	0.02785949276	0.01851259253	0.01312895682	0.00978483286	0.00756521107
26	0.04290579437	0.02769531699	0.01844841522	0.01311148673	0.00978017739	0.00756388405
28	0.04280033153	0.02756430324	0.01840481529	0.01310126617	0.00977778854	0.00756328233
30	0.04270993496	0.02745937089	0.01837513880	0.01309525214	0.00977655328	0.00756300336
extra.	0.04170186316	0.02715342577	0.01832502644	0.01302827108	0.00976919713	0.00755456686
exact	0.04171854326	0.02702161935	0.01831051051	0.01308654379	0.00977523196	0.00756276973

and $\langle S_j^x S_{j+2}^x \rangle$:

$$P(3) = \frac{1}{2} + \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x} \frac{1}{8} \left[\frac{x}{\sin^2(\phi x)} \left(\frac{3 \sinh^2 \phi}{\sin^2(\phi x)} - 5 - \cosh 2\phi \right) - \cot(\phi x) \left(\frac{3 \tanh \phi}{\sin^2(\phi x)} - \frac{4 + 8 \cosh 2\phi}{\sinh 2\phi} \right) \right], \quad (36)$$

$$\langle S_j^x S_{j+2}^x \rangle = \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x} \frac{1}{8} \left[-\frac{x}{\sin^2(\phi x)} \left(\frac{3 \sinh^2 \phi}{\sin^2(\phi x)} + 1 - 3 \cosh 2\phi \right) + \cot(\phi x) \left(\frac{3 \cosh(2\phi) \tanh(\phi)}{\sin^2(\phi x)} - \frac{4}{\sinh(2\phi)} \right) \right]. \quad (37)$$

Using equations (28, 29) we have

$$\langle S_j^z S_{j+1}^z \rangle = \frac{1}{4} + \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x} \frac{1}{2} \left(\cot(\phi x) \coth(\phi) - \frac{x}{\sin^2 \phi x} \right), \quad (38)$$

$$\langle S_j^z S_{j+2}^z \rangle = \frac{1}{4} + \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x} \frac{1}{4} \left[\frac{x}{\sin^2(\phi x)} \left(\frac{3 \sinh^2 \phi}{\sin^2(\phi x)} - 1 - \cosh 2\phi \right) - \cot(\phi x) \left(\frac{3 \tanh \phi}{\sin^2(\phi x)} - 4 \coth 2\phi \right) \right]. \quad (39)$$

IV. NUMERICAL RESULTS AND CONCLUSION

We calculate numerically the second neighbor correlations $\langle S_0^z S_2^z \rangle$ and $\langle S_0^x S_2^x \rangle$ for finite rings with $N = 20, 22, 24, 26, 28, 30$. In Table (I) and (II) results of (39) and (37) are compared with numerical results. They coincide very well.

In the previous paper we treated the second neighbor correlations for massless region $1 > \Delta > -1$. In this paper we have shown that almost the same method can be applied for massive region $\Delta > 1$. This is the generalization of $\Delta = 1$ result

$$\langle S_j^z S_{j+2}^z \rangle = \frac{1}{12} - \frac{4}{3} \ln 2 + \frac{3}{4} \zeta(3). \quad (40)$$

The multiple integrals for the correlation functions of the

XXZ chain at the massive region can be reduced to the one-dimensional ones in the case of the next nearest-neighbor correlation functions. This property will be generalized to the third neighbor or longer distance correlations. In our future work, we are particularly interested in calculating the third-neighbor correlation functions $\langle S_j^x S_{j+3}^x \rangle$ and $\langle S_j^z S_{j+3}^z \rangle$ for general Δ . For $\Delta = 1$, they were recently calculated in [13]

$$\langle S_j^z S_{j+3}^z \rangle = \frac{1}{12} - 3 \ln 2 + \frac{37}{6} \zeta(3) - \frac{14}{3} \zeta(3) \ln 2 - \frac{3}{2} \zeta(3)^2 - \frac{125}{24} \zeta(5) + \frac{25}{3} \zeta(5) \ln 2. \quad (41)$$

from the multiple integrals. This should be generalized to arbitrary Δ .




FIG. 1: Nearest-neighbor and next nearest neighbor correlation functions for the XXZ chain. We calculated $\langle S_j^z S_{j+1}^z \rangle$, $\langle S_j^x S_{j+1}^x \rangle$, $\langle S_j^z S_{j+2}^z \rangle$ and $\langle S_j^x S_{j+2}^x \rangle$

Acknowledgments

We are grateful to K. Sakai for valuable discussions. The numerical data for finite rings have been obtained by the Fortran package TITPACK Ver. 2. This work

is in part supported by Grant-in-Aid for the Scientific Research (B) No. 14340099 from the Ministry of Education, Culture, Sports, Science and Technology, Japan. One of the authors (GK) is supported by the JSPS research fellowships for young scientists. MS is supported by Grant-in Aid for Young Scientists (B) No. 14740228.

APPENDIX A: ANTISYMMETRIC DIFFERENTIAL OPERATOR

Elliptic theta function satisfies the following quasi double periodicity

$$\vartheta_1(\lambda + \pi) = -\vartheta_1(\lambda), \quad \vartheta_1(\lambda + i\phi) = -e^\phi e^{-2i\lambda} \vartheta_1(\lambda). \quad (\text{A1})$$

We consider

$$f(\lambda_1, \lambda_2) = \left(\frac{\partial}{\partial \lambda_2} - \frac{\partial}{\partial \lambda_1} \right) U_1(\lambda_1) U_1(\lambda_2).$$

Using (A1) and $\vartheta_2(\lambda) = \vartheta_1(\lambda + \pi/2)$ we can show that $U_1(\lambda)$ and $f(\lambda_1, \lambda_2)$ are elliptic functions which satisfy

$$\begin{aligned} U_1(\lambda_2) &= -U_1(\lambda_2 + i\phi) = U_1(\lambda_2 + \pi), \\ f(\lambda_1, \lambda_2) &= -f(\lambda_1, \lambda_2 + i\phi) = f(\lambda_1, \lambda_2 + \pi). \end{aligned} \quad (\text{A2})$$

As $f(\lambda_1, \lambda_2) = -f(\lambda_2, \lambda_1)$, f has zeros at $\lambda_1 + k\pi + il\phi$ as function of λ_2 . It has poles at $k\pi + i(l - 1/2)\phi$ with order 2. Then function f should be factorized as follows

$$f(\lambda_2) = C(\lambda_1) \frac{\vartheta_1(\lambda_2 - \lambda_1) \vartheta_1(\lambda_2 - \alpha)}{\vartheta_1(\lambda_2 + i\phi/2)^2}. \quad (\text{A3})$$

Here α is another zero. Using (A1) we have $f(\lambda_2 + i\phi) = \exp(2i(\alpha + \lambda_1 + i\phi)) f(\lambda_2)$. By the condition (A2) α must be $\pi/2 - \lambda_1 - i\phi$. The residue of $U_1(\lambda)$ at $\lambda = -i\phi/2$ is i . Then we have

$$\lim_{\lambda_2 \rightarrow -i\phi/2} (\lambda_2 + i\phi/2)^2 f(\lambda_1, \lambda_2) = -iU_1(\lambda_1).$$

By this condition we find

$$C(\lambda_1) = iU_1(\lambda_1) \frac{(\vartheta_1'(0))^2}{\vartheta_1(\lambda_1 + i\phi/2) \vartheta_1(\lambda_1 + i\phi/2 + \pi/2)}, \quad (\text{A4})$$

and therefore $f(\lambda_1, \lambda_2) = U_2(\lambda_1, \lambda_2)$. Thus we have proven (20) for $m = 2$.

To prove for $m \geq 3$ we assume that it stands for U_{m-1} . Define

$$f_m(\lambda_1, \dots, \lambda_m) = \frac{1}{(m-1)!} \prod_{l=1}^{m-1} \left(\frac{\partial}{\partial \lambda_m} - \frac{\partial}{\partial \lambda_l} \right) U_1(\lambda_m) U_{m-1}(\lambda_1, \dots, \lambda_{m-1}). \quad (\text{A5})$$

This function has the following properties:

$$f_m(\lambda_1, \dots, \lambda_m) = -f_m(\lambda_1, \dots, \lambda_m + i\phi) = f_m(\lambda_1, \dots, \lambda_m + \pi). \quad (\text{A6})$$

As f_m is antisymmetric for exchange of variables we know $m - 1$ zeroes and m fold pole:

$$f_m(\lambda_m) = C(\lambda_1, \dots, \lambda_{m-1}) \frac{\prod_{l=1}^{m-1} \vartheta_1(\lambda_m - \lambda_l) \vartheta_1(\lambda_m - \alpha)}{\vartheta_1(\lambda_m + i\phi/2)^m}. \quad (\text{A7})$$

α must be $\pi/2 - \sum_{l=1}^{m-1} \lambda_l - im\phi/2$ by the condition (A6). Using

$$\lim_{\lambda_m \rightarrow -i\phi/2} (\lambda_m + i\frac{\phi}{2})^m f_m(\lambda_1, \dots, \lambda_m) = (-1)^{m-1} i U_{m-1}(\lambda_1, \dots, \lambda_{m-1}),$$

we can determine C ,

$$C(\lambda_1, \dots, \lambda_{m-1}) = i U_{m-1}(\lambda_1, \dots, \lambda_{m-1}) \frac{(\vartheta'_1(0))^m}{\vartheta_1(\sum_{l=1}^{m-1} (\lambda_l + i\phi/2) + \pi/2) \prod_{l=1}^{m-1} \vartheta_1(\lambda_l + i\phi/2)}. \quad (\text{A8})$$

Thus we show

$$U_m(\lambda_1, \dots, \lambda_m) = \frac{1}{\prod_{r=1}^{m-1} r!} \prod_{j>k} \left(\frac{\partial}{\partial \lambda_j} - \frac{\partial}{\partial \lambda_k} \right) U_1(\lambda_1) U_1(\lambda_2) \dots U_1(\lambda_m). \quad (\text{A9})$$

for general m stands if it stands for $m - 1$. Putting $\lambda_j = \phi x_j$ we obtain (20).

APPENDIX B: SOME INTEGRAL FORMULA

For the integration of canonical forms following formula is important,

$$\begin{aligned} \int_{-Q/2}^{Q/2} dx_1 e^{2in\phi x_1} \mathbf{W}_1(x_1 + \epsilon_1) &= \int_{-Q/2}^{Q/2} dx_1 e^{2in\phi x_1 - \epsilon_1} \mathbf{W}_1(x_1) \\ &= \int_{-\infty}^{\infty} dx_1 e^{2in\phi(x_1 - \epsilon_1)} \frac{1}{2} \text{sech}(\pi x_1) = \frac{\exp(-2in\phi\epsilon_1)}{2 \cosh n\phi}. \end{aligned} \quad (\text{B1})$$

Here we have used (19). Next we consider the integral of $e^{2in\phi x_1} / (e^{2i\phi x_2} - e^{2i\phi x_1})$:

$$\begin{aligned} \int_{-Q/2}^{Q/2} dx_1 dx_2 \frac{e^{2in\phi x_1}}{e^{2i\phi x_2} - e^{2i\phi x_1}} \mathbf{W}_1(x_1 + \epsilon_1) \mathbf{W}_1(x_2 + \epsilon_2) \\ = -i \int_{-Q/2}^{Q/2} \int_{-Q/2}^{Q/2} \frac{\exp(i\phi((2n-1)x_1 - x_2))}{2 \sin \phi(x_2 - x_1)} \mathbf{W}_1(x_1 + \epsilon_1) \mathbf{W}_1(x_2 + \epsilon_2) dx_1 dx_2 \\ = -ie^{i\phi(-(2n-1)\epsilon_1 + \epsilon_2)} \int_{-Q/2}^{Q/2} \int_{-Q/2}^{Q/2} \frac{\exp(i\phi((2n-1)x_1 - x_2))}{2 \sin \phi(x_2 - x_1 + \epsilon_1 - \epsilon_2)} \mathbf{W}_1(x_1) \mathbf{W}_1(x_2) dx_1 dx_2 \end{aligned}$$

Using (19) we extend the region of integration:

$$= -ie^{i\phi(-(2n-1)\epsilon_1 + \epsilon_2)} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{\exp(i\phi((2n-1)x_1 - x_2))}{8 \sin \phi(x_2 - x_1 + \epsilon_1 - \epsilon_2)} \text{sech} \pi x_1 \text{sech} \pi x_2 dx_1 dx_2.$$

The term $\text{sech} \pi x_1 \text{sech} \pi x_2$ can be written as $(\tanh \pi x_2 - \tanh \pi x_1) / \sinh \pi(x_2 - x_1)$. Putting $y_1 = x_1 + x_2, y_2 = x_2 - x_1$ we have

$$= -ie^{i\phi(-(2n-1)\epsilon_1 + \epsilon_2)} \int_{-\infty}^{\infty} \frac{dy_2 \exp(-i\phi n y_2)}{16 \sinh \pi y_2 \sin \phi(y_2 + \epsilon_1 - \epsilon_2)} \int_{-\infty}^{\infty} e^{i(n-1)\phi y_1} (\tanh \frac{\pi}{2}(y_1 + y_2) - \tanh \frac{\pi}{2}(y_1 - y_2)) dy_1.$$

For $n \neq 1$ the last integral is estimated as follows:

$$\begin{aligned}
& \int_{-\infty}^{\infty} e^{i(n-1)\phi y_1} \left(\tanh \frac{\pi}{2}(y_1 + y_2) - \tanh \frac{\pi}{2}(y_1 - y_2) \right) dy_1 \\
&= -\frac{\pi}{2} \int_{-\infty}^{\infty} \frac{e^{i(n-1)\phi y_1} - 1}{(n-1)\phi i} \left(\frac{1}{\cosh^2 \frac{\pi}{2}(y_1 + y_2)} - \frac{1}{\cosh^2 \frac{\pi}{2}(y_1 - y_2)} \right) dy_1 \\
&= \frac{\pi}{2} \int_{-\infty}^{\infty} \frac{e^{i(n-1)\phi(y_1+y_2)} - e^{i(n-1)\phi(y_1-y_2)}}{(n-1)\phi i} \frac{dy_1}{\cosh^2 \frac{\pi}{2} y_1} = \pi \frac{\sin(n-1)\phi y_2}{(n-1)\phi} \int_{-\infty}^{\infty} e^{i\phi(n-1)y_1} \frac{dy_1}{\cosh^2 \frac{\pi}{2} y_1} \\
&= \frac{4 \sin(n-1)\phi y_2}{\sinh(n-1)\phi}.
\end{aligned}$$

For $n = 1$ this is $4y_2$. Then only integral with respect to y_2 remains. To circumvent the poles on the real axis we change the path of integral from $[-\infty, \infty]$ to $[-\infty + i/2, \infty + i/2]$. Thus we have

$$\begin{aligned}
& \int_{-Q/2}^{Q/2} dx_1 dx_2 \frac{e^{2in\phi x_1}}{e^{2i\phi x_2} - e^{2i\phi x_1}} \mathbf{W}_1(x_1 + \epsilon_1) \mathbf{W}_1(x_2 + \epsilon_2) \\
&= \begin{cases} -ie^{i\phi(-(2n-1)\epsilon_1+\epsilon_2)} \int_{-\infty+i/2}^{\infty+i/2} \frac{dx \exp(-ni\phi x)}{4 \sinh \pi x \sin \phi(x + \epsilon_1 - \epsilon_2)} \frac{\sin(n-1)\phi x}{\sinh(n-1)\phi}, & \text{for } n \neq 1, \\ -ie^{i\phi(-\epsilon_1+\epsilon_2)} \int_{-\infty+i/2}^{\infty+i/2} \frac{dx \exp(-i\phi x)}{4 \sinh \pi x \sin \phi(x + \epsilon_1 - \epsilon_2)} x, & \text{for } n = 1. \end{cases} \quad (\text{B2})
\end{aligned}$$

If the integrand is given by a canonical form we can estimate the integral by these formulae.

APPENDIX C: CANONICAL FORM OF INTEGRAND

The weight function $\mathbf{W}_m$ is elliptic function with periodicity $Q, 2i$ and the integrand is periodic function with periodicity Q . Then integrand is expressed by $z_l \equiv \exp(2i\phi x_l + \phi)$. If integrand has the form $(z_1 - 1)^m f(z_1, z_2, \dots, z_m)$ and f has no singularity as a function of z_1 , the m -th pole of $\mathbf{W}_m$ at $x_1 = i/2$ is cancelled and shift of path from $[-Q/2, Q/2]$ to $[-Q/2 + i, Q/2 + i]$ becomes possible:

$$\int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m(z_1 - 1)^m f(z_1, z_2, \dots) = - \int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m(q^2 z_1 - 1)^m f(q^2 z_1, z_2, \dots). \quad (\text{C1})$$

Here we have used (23). In the same way we have

$$\int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m(q^2 z_1 - 1)^m f(z_1, z_2, \dots) = - \int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m(z_1 - 1)^m f(q^{-2} z_1, z_2, \dots). \quad (\text{C2})$$

Actually f might have single pole at $z_1 = q^{2n} z_2$, $n = \text{integer}$, because $\mathbf{W}_m$ has single zeros at these points. Then we have

$$\int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m \frac{(z_1 - 1)^m}{z_1 - q^2 z_2} f(z_1, z_2, \dots) = - \int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m \frac{(q^2 z_1 - 1)^m}{q^2(z_1 - z_2)} f(q^2 z_1, z_2, \dots), \quad (\text{C3})$$

$$\int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m \frac{(q^2 z_1 - 1)^m}{q^2 z_1 - z_2} f(z_1, z_2, \dots) = - \int_{-Q/2}^{Q/2} dx_1 \mathbf{W}_m \frac{(z_1 - 1)^m}{z_1 - z_2} f(q^{-2} z_1, z_2, \dots). \quad (\text{C4})$$

From (22) we have

$$\int_{-Q/2}^{Q/2} d^m x \mathbf{W}_m g(z_1, z_2, z_3, \dots) = - \int_{-Q/2}^{Q/2} d^m x \mathbf{W}_m g(z_2, z_1, z_3, \dots). \quad (\text{C5})$$

Suffix 1 and 2 can be replaced by general $j \neq k$ at $1 \leq j, k \leq m$. These rules are sufficient to simplify the integrand. Hereafter we write $A \sim B$ if

$$\int_{-Q/2}^{Q/2} d^m x \mathbf{W}_m A = \int_{-Q/2}^{Q/2} d^m x \mathbf{W}_m B.$$

At first we consider two-dimensional integral. Equation (30) and (31) are rewritten in terms of $q \equiv \exp(-\phi)$ and $z_l \equiv \exp(2i\phi x_l + \phi)$,

$$P(2) = \int_{-Q/2}^{Q/2} dx_1 dx_2 \mathbf{W}_2 \frac{(q^2 z_1 - 1)(z_2 - 1)}{2i\phi(z_2 - q^2 z_1)}. \quad (\text{C6})$$

The integrand is transformed as follows

$$\frac{(q^2 z_1 - 1)(z_2 - 1)}{2i\phi(z_2 - q^2 z_1)} = \frac{1}{2i\phi} \left(-z_2 + 1 + \frac{(z_2 - 1)^2}{z_2 - q^2 z_1} \right) \sim \frac{1}{2i\phi} \left(-z_2 - \frac{(q^2 z_2 - 1)^2}{q^2(z_2 - z_1)} \right). \quad (\text{C7})$$

We can show that $z_1^2/(z_2 - z_1) \sim -z_1 + q^{-2}/(z_2 - z_1)$, because

$$\begin{aligned} \frac{(z_1 - 1)^2}{z_2 - z_1} &\sim -\frac{(q^2 z_1 - 1)^2}{z_2 - q^2 z_1} = q^2 z_1 + z_2 - 2 - \frac{(z_2 - 1)^2}{z_2 - q^2 z_1} \\ &\sim -(1 - q^2)z_1 + \frac{(q^2 z_2 - 1)^2}{q^2(z_2 - z_1)} \sim -(1 - q^2)z_1 + \frac{(q^2 z_1 - 1)^2}{q^2(z_2 - z_1)}. \end{aligned} \quad (\text{C8})$$

Then we have the canonical form

$$(\text{C7}) \sim \frac{1}{2i\phi} \left((1 + q^2)z_1 + \frac{2z_1 - 1 - q^{-2}}{z_2 - z_1} \right). \quad (\text{C9})$$

Using (B1) and (B2) we evaluate $\int_{-Q/2}^{Q/2} dx_1 dx_2 \mathbf{W}_1(x_1 + \epsilon_1) \mathbf{W}_1(x_2 + \epsilon_2)$ (C9):

$$\frac{e^{-2i\phi\epsilon_1}}{4i\phi} + \frac{1}{4\phi} \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x \sin \phi(x + \epsilon_1 - \epsilon_2)} \left[-x e^{-i\phi(x+\epsilon_1-\epsilon_2)} + e^{i\phi(\epsilon_1+\epsilon_2)} \coth \phi \sin \phi x \right]. \quad (\text{C10})$$

Using (20) we have

$$P(2) = \lim_{\epsilon_1, \epsilon_2 \rightarrow 0} (\partial_2 - \partial_1)(\text{C10}) = (34). \quad (\text{C11})$$

(31) is rewritten

$$\langle S_j^x S_{j+1}^x \rangle = \int_{-Q/2}^{Q/2} dx_1 dx_2 \mathbf{W}_2 \frac{(q^2 z_1 - 1)(z_2 - 1)}{4i\phi q(z_2 - z_1)}. \quad (\text{C12})$$

The canonical form of integrand is

$$\frac{1}{4i\phi q} \frac{-(1 + q^2)z_1 + 2}{z_2 - z_1}. \quad (\text{C13})$$

Then we have

$$\langle S_j^x S_{j+1}^x \rangle = \lim_{\epsilon_1, \epsilon_2 \rightarrow 0} (\partial_2 - \partial_1) \frac{1}{8\phi} \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x \sin \phi(x + \epsilon_1 - \epsilon_2)} \left[x \cosh \phi e^{-i\phi(x+\epsilon_1-\epsilon_2)} - e^{i\phi(\epsilon_1+\epsilon_2)} \frac{\sin \phi x}{\sinh \phi} \right] = (35). \quad (\text{C14})$$

Equation (32) is

$$P(3) = \frac{i}{8\phi^3} \int_{-Q/2}^{Q/2} d^3 x \mathbf{W}_3 \frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{(z_2 - q^2 z_1)(z_3 - q^2 z_1)(z_3 - q^2 z_2)}, \quad (\text{C15})$$

The integrand is devided as follows

$$\begin{aligned} \frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{(z_2 - q^2 z_1)(z_3 - q^2 z_1)(z_3 - q^2 z_2)} &= \frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{z_2(1 - q^2)} \\ &\times \left(\frac{1}{(z_3 - q^2 z_1)(z_3 - q^2 z_2)} + \frac{1}{(z_3 - q^2 z_1)(z_2 - q^2 z_1)} - \frac{1}{(z_3 - q^2 z_2)(z_2 - q^2 z_1)} \right) \end{aligned}$$

The first and second terms are symmetrized by the exchange of variables and the third term is divided to four terms

$$\begin{aligned} & \sim -(q^2 z_1 - 1)(q^2 z_2 - 1) \frac{(z_3 - 1)^2(1 + q^2(z_1 z_2 - z_1 - z_2))}{q^2 z_1 z_2 (1 - q^2)(z_3 - q^2 z_2)} \\ & + (z_1 - 1)(z_2 - 1) \frac{(q^2 z_1 - 1)^2(1 + q^2 z_2 z_3 - z_2 - z_3)}{z_2 z_3 (1 - q^2)(z_2 - q^2 z_1)} + \frac{(q^2 z_1 - 1)^2(z_3 - 1)^2}{z_2(1 - q^2)} \\ & - \frac{(q^2 z_1 - 1)^2(z_3 - 1)^3}{z_2(1 - q^2)(z_3 - q^2 z_2)} + \frac{(z_3 - 1)^2(q^2 z_1 - 1)^3}{z_2(1 - q^2)(z_2 - q^2 z_1)} - \frac{(z_3 - 1)^3(q^2 z_1 - 1)^3}{z_2(1 - q^2)(z_3 - q^2 z_2)(z_2 - q^2 z_1)}. \end{aligned} \quad (C16)$$

This is equivalent to

$$\begin{aligned} & \sim \left(\frac{(q^2 z_1 - 1)^2(z_3 - 1)^2}{z_2(1 - q^2)} - \frac{(q^2 z_1 - 1)^2(z_3 - 1)^2 - (q^2 z_3 - 1)^2(z_1 - 1)^2 q^2}{q^2 z_1 z_2 (1 - q^2)} \right. \\ & + \frac{(q^2 z_1 - 1)(z_1 - 1)((z_3 - 1)^2 - (q^2 z_3 - 1)^2)}{z_1(1 - q^2)} + \frac{(q^2 z_3 - 1)(z_3 - 1)(q^2 z_1 - 1)^3 - (z_2 - 1)^3}{z_3(1 - q^2)(z_2 - q^2 z_1)} \Big) \\ & + \left(-\frac{(q^2 z_1 - 1)^3(z_3 - 1)^3}{q^2 z_1 z_2 (1 - q^2)(z_3 - q^2 z_2)} + \frac{(z_3 - 1)^3(q^2 z_1 - 1)^3}{z_2 z_3 (1 - q^2)(z_2 - q^2 z_1)} - \frac{(z_3 - 1)^3(q^2 z_1 - 1)^3}{z_2(1 - q^2)(z_3 - q^2 z_2)(z_2 - q^2 z_1)} \right). \end{aligned} \quad (C17)$$

We use following formula to simplify terms in the second parenthesis:

$$\frac{(z_3 - 1)^3}{z_3 - q^2 z_2} f(z_1, z_2) \sim -\frac{(q^2 z_3 - 1)^3}{q^2(z_3 - z_2)} f(z_1, z_2), \quad \frac{(q^2 z_1 - 1)^3}{z_2 - q^2 z_1} f(z_2, z_3) \sim -\frac{(z_1 - 1)^3}{q^2(z_2 - z_1)} f(z_2, z_3). \quad (C18)$$

The first parenthesis is simplified only by the exchange of variables. Then we have

$$\begin{aligned} (C17) & \sim \left(\frac{3q^2 z_1^2 z_3}{z_2} - 3(1 + q^2) \frac{z_1^2}{z_2} + 8 \frac{z_1}{z_2} \right) + \left(\frac{(q^2 z_3 - 1)^3(q^2 z_1 - 1)^3 - q^4(z_3 - 1)^3(z_1 - 1)^3}{q^4(1 - q^2)z_2 z_3(z_2 - z_1)} \right. \\ & \left. - \frac{(q^2 z_3 - 1)^3(z_1 - 1)^3}{z_2 q^2(1 - q^2)(z_3 - z_2)(z_2 - z_1)} \right). \end{aligned} \quad (C19)$$

The last term in the second parenthesis is

$$\sim -\frac{(q^2 z_3 - 1)^3(z_1 - 1)^3 - (q^2 z_1 - 1)^3(z_3 - 1)^3}{q^2(1 - q^2)z_2(z_2 - z_1)}.$$

The terms of the form $f(z_1, z_3)/(z_2(z_2 - z_1))$ can be replaced by $-f(z_1, z_3)/(z_1 z_2) + f(z_1, z_3)/(z_1(z_2 - z_1))$. Then we have

$$\begin{aligned} (C19) & \sim \left(\frac{3q^2 z_1^2 z_3}{z_2} - 3(1 + q^2) \frac{z_1^2}{z_2} + 8 \frac{z_1}{z_2} \right) - \frac{(q^2 z_3 - 1)^3(q^2 z_1 - 1)^3 - q^4(z_3 - 1)^3(z_1 - 1)^3}{q^4(1 - q^2)z_1 z_2 z_3} \\ & + \frac{(q^2 z_3 - 1)^2(z_1 - 1)^2 + (q^2 z_3 - 1)(z_3 - 1)(q^2 z_1 - 1)(z_1 - 1) + (q^2 z_1 - 1)^2(z_3 - 1)^2}{q^2 z_1 z_2} \\ & + \frac{1}{z_2 - z_1} \left((q^6 + q^4 + q^2 + 1)z_1^2 z_3^2 - 3(q^4 + q^2 + 1)z_1 z_3(z_1 + z_3) + 3(q^2 + 1)(z_1^2 + 3z_1 z_3 + z_3^2) \right. \\ & \left. - \left(\frac{z_1^2}{z_3} + \frac{z_3^2}{z_1} + 9(z_1 + z_3) \right) + \frac{3}{q^2} \left(\frac{1}{z_1} + \frac{1}{z_3} \right) - \frac{1 + q^{-4}}{z_1 z_3} \right. \\ & \left. - \frac{(q^2 z_3 - 1)^2(z_1 - 1)^2 + (q^2 z_3 - 1)(z_3 - 1)(q^2 z_1 - 1)(z_1 - 1) + (q^2 z_1 - 1)^2(z_3 - 1)^2}{q^2 z_1} \right). \end{aligned} \quad (C20)$$

Then all terms are in the form of $z_1^a z_2^b z_3^c$ or $z_1^a z_3^b/(z_2 - z_1)$ with $2 \geq a, b, c \geq -1$. We should note the following relations

$$\begin{aligned} z_1^2 f(z_2, z_3) & \sim \left(\frac{3(1 + q^2)}{1 + q^4} z_1 - \frac{6}{1 + q^4} + \frac{1 + q^2}{q^2(1 + q^4)z_1} \right) f(z_2, z_3), \\ \frac{z_1^2}{z_2 - z_1} f(z_3) & \sim \left(\frac{3}{q^2(z_2 - z_1)} - \frac{1 + q^2}{q^4(z_2 - z_1)z_1} - z_1 \right) f(z_3). \end{aligned} \quad (C21)$$

By these relations we can put powers a, b, c are at 1, 0, -1. The canonical form is

$$(C20) \sim \frac{(1+q^2)^2}{q^2} \frac{z_1}{z_3} + \frac{1}{(z_2 - z_1)} \left[\frac{12(1+q^2)}{q^2} - \frac{(1+10q^2+q^4)z_1}{q^2} - \frac{3(1+q^2)^2}{q^4 z_1} \right. \\ \left. + z_3 \left\{ -\frac{1+10q^2+q^4}{q^2} + 3(1+q^2)z_1 + \frac{3(1+q^2)}{q^2 z_1} \right\} + \frac{1}{z_3} \left\{ -\frac{3(1+q^2)^2}{q^4} + \frac{3(1+q^2)z_1}{q^2} + \frac{(1+q^2)(1+q^2+q^4)}{q^6 z_1} \right\} \right]. \quad (C22)$$

Multiple integral can be done through (B1, B2);

$$\frac{i}{8\phi^3} \int_{-Q/2}^{Q/2} d^3x \mathbf{W}_1(x_1 + \epsilon_1) \mathbf{W}_1(x_2 + \epsilon_2) \mathbf{W}_1(x_3 + \epsilon_3) (C22) = i \frac{e^{2i(\epsilon_3 - \epsilon_1)\phi}}{16\phi^3} + \frac{1}{8\phi^3} \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x \sin \phi(x + \epsilon_1 - \epsilon_2)} \\ \times \left[\frac{12(1+q^2)}{q} e^{i\phi(\epsilon_1 + \epsilon_2)} \frac{\sin \phi x}{\sinh \phi} - \frac{(1+10q^2+q^4)}{q^2} e^{i\phi(-x - \epsilon_1 + \epsilon_2)} x - \frac{3(1+q^2)^2}{q^2} e^{i\phi(x + 3\epsilon_1 + \epsilon_2)} \frac{\sin 2\phi x}{\sinh 2\phi} \right. \\ \left. + \frac{e^{-2i\phi\epsilon_3}}{\cosh \phi} \left\{ -\frac{1+10q^2+q^4}{q^2} e^{i\phi(\epsilon_1 + \epsilon_2)} \frac{\sin \phi x}{\sinh \phi} + \frac{3(1+q^2)}{q} e^{i\phi(-x - \epsilon_1 + \epsilon_2)} x + \frac{3(1+q^2)}{q} e^{i\phi(x + 3\epsilon_1 + \epsilon_2)} \frac{\sin 2\phi x}{\sinh 2\phi} \right\} \right. \\ \left. + \frac{e^{2i\phi\epsilon_3}}{\cosh \phi} \left\{ -\frac{3(1+q^2)^2}{q^2} e^{i\phi(\epsilon_1 + \epsilon_2)} \frac{\sin \phi x}{\sinh \phi} + \frac{3(1+q^2)}{q} e^{i\phi(-x - \epsilon_1 + \epsilon_2)} x + \frac{(1+q^2)(1+q^2+q^4)}{q^3} e^{i\phi(x + 3\epsilon_1 + \epsilon_2)} \frac{\sin 2\phi x}{\sinh 2\phi} \right\} \right]. \quad (C23)$$

Then we have

$$P(3) = \frac{1}{2} \lim_{\epsilon_1, \epsilon_2, \epsilon_3 \rightarrow 0} (\partial_2 - \partial_1)(\partial_3 - \partial_1)(\partial_3 - \partial_2)(C23) = (36). \quad (C24)$$

Equation (33) is

$$\langle S_j^x S_{j+2}^x \rangle = \frac{i}{8\phi^3} \int_{-Q/2}^{Q/2} d^3x \mathbf{W}_3 \frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{q^2 (z_2 - z_1)(z_3 - z_1)(z_3 - q^2 z_2)}. \quad (C25)$$

The integrand is devided as follows

$$\frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{q^2 (1 - q^2) z_2} \left(\frac{1}{(z_3 - z_1)(z_3 - q^2 z_2)} + \frac{1}{(z_3 - z_1)(z_2 - z_1)} - \frac{1}{(z_3 - q^2 z_2)(z_2 - z_1)} \right). \quad (C26)$$

The first term is transformed as

$$\frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{q^2 (1 - q^2) z_2 (z_3 - z_1)(z_3 - q^2 z_2)} = \frac{(q^2 z_1 - 1)^2 (z_2 - 1)(q^2 z_2 - 1)}{q^2 (1 - q^2) z_2 (z_3 - z_1)} (z_3 + q^2 z_2 - 2 + \frac{(q^2 z_2 - 1)^2}{z_3 - q^2 z_2}) \\ \sim \frac{(q^2 z_1 - 1)^2 (z_2 - 1)(q^2 z_2 - 1)}{q^2 (1 - q^2) z_2 (z_3 - z_1)} (z_3 + q^2 z_2 - 2) - \frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)^3}{(1 - q^2) z_2 (z_3 - z_1)(z_3 - z_2)}. \quad (C27)$$

The third term is transformed as

$$- \frac{(q^2 z_1 - 1)^2 (q^2 z_2 - 1)(z_2 - 1)(z_3 - 1)^2}{q^2 (1 - q^2) z_2 (z_3 - q^2 z_2)(z_2 - z_1)} = - \frac{(q^2 z_1 - 1)^2 (z_2 - 1)(z_3 - 1)^2}{q^2 (1 - q^2) z_2 (z_2 - z_1)} \left(-1 + \frac{z_3 - 1}{z_3 - q^2 z_2} \right) \\ \sim \frac{(q^2 z_1 - 1)^2 (z_2 - 1)(z_3 - 1)^2}{q^2 (1 - q^2) z_2 (z_2 - z_1)} + \frac{(q^2 z_1 - 1)^2 (z_2 - 1)(q^2 z_3 - 1)^3}{q^4 (1 - q^2) z_2 (z_2 - z_1)(z_3 - z_2)}. \quad (C28)$$

Using the equivalence

$$\frac{f(z_1, z_2, z_3)}{(z_1 - z_2)(z_1 - z_3)} \sim \frac{1}{z_1 - z_2} \left(\frac{f(z_1, z_2, z_3) - f(z_1, z_3, z_2)}{z_2 - z_3} \right), \quad (C29)$$

we replace the term like $(z_1 - z_2)(z_1 - z_3)$ in the denominator by $z_1 - z_2$. The canonical form is

$$(C26) \sim \frac{1}{z_2 - z_1} \left[-\frac{(1+q^2)(3+q^4)}{q^4} + \frac{(3-2q^2+3q^4)z_1}{q^2} + \frac{(1+q^2)^2}{q^6 z_1} \right. \\ \left. + z_3 \left\{ \frac{6-2q^2}{q^2} - \frac{(1+q^2)z_1}{q^2} + \frac{(1+q^2)(-2+q^2)}{q^4 z_1} \right\} + \frac{1}{z_3} \left\{ \frac{4+4q^2}{q^2} - (1+q^2)z_1 - \frac{1+q^2}{q^4 z_1} \right\} \right]. \quad (C30)$$

Then the integral of this form is

$$\begin{aligned}
& \frac{i}{8\phi^3} \int_{-Q/2}^{Q/2} d^3x \mathbf{W}_1(x_1 + \epsilon_1) \mathbf{W}_1(x_2 + \epsilon_2) \mathbf{W}_1(x_3 + \epsilon_3) \quad (\text{C30}) \\
&= \frac{1}{8\phi^3} \int_{-\infty+i/2}^{\infty+i/2} \frac{dx}{\sinh \pi x \sin \phi(x + \epsilon_1 - \epsilon_2)} \\
&\times \left[-\frac{(1+q^2)(3+q^4)}{q^3} e^{i\phi(\epsilon_1+\epsilon_2)} \frac{\sin \phi x}{\sinh \phi} + \frac{(3-2q^2+3q^4)}{q^2} e^{i\phi(-x-\epsilon_1+\epsilon_2)} x + \frac{(1+q^2)^2}{q^2} e^{i\phi(x+3\epsilon_1+\epsilon_2)} \frac{\sin 2\phi x}{\sinh 2\phi} \right. \\
&+ \frac{e^{-2i\phi\epsilon_3}}{\cosh \phi} \left\{ \frac{6-2q^2}{q} e^{i\phi(\epsilon_1+\epsilon_2)} \frac{\sin \phi x}{\sinh \phi} - \frac{(1+q^2)}{q} e^{i\phi(-x-\epsilon_1+\epsilon_2)} x + \frac{(1+q^2)(-2+q^2)}{q^2} e^{i\phi(x+3\epsilon_1+\epsilon_2)} \frac{\sin 2\phi x}{\sinh 2\phi} \right\} \\
&+ \left. \frac{e^{2i\phi\epsilon_3}}{\cosh \phi} \left\{ \frac{4+4q^2}{q} e^{i\phi(\epsilon_1+\epsilon_2)} \frac{\sin \phi x}{\sinh \phi} - \frac{1+q^2}{q} e^{i\phi(-x-\epsilon_1+\epsilon_2)} x - \frac{1+q^2}{q} e^{i\phi(x+3\epsilon_1+\epsilon_2)} \frac{\sin 2\phi x}{\sinh 2\phi} \right\} \right]. \quad (\text{C31})
\end{aligned}$$

Then we have

$$\langle S_j^x S_{j+2}^x \rangle = \frac{1}{2} \lim_{\epsilon_1, \epsilon_2, \epsilon_3 \rightarrow 0} (\partial_2 - \partial_1)(\partial_3 - \partial_1)(\partial_3 - \partial_2) (\text{C31}) = (37). \quad (\text{C32})$$

-
- | | |
|--|---|
| <p>[1] H.A. Bethe, Z. Phys. 71 (1931) 205.
 [2] L. Hulthén, Ark. Mat. Astron. Fys. A 26 (1938) 1.
 [3] M. Takahashi, <i>Thermodynamics of One-Dimensional Solvable Models</i>, (Cambridge University Press, 1999).
 [4] M. Takahashi, J. Phys. C 10 (1977) 1289.
 [5] M. Jimbo, K. Miki, T. Miwa and A. Nakayashiki, Phys. Lett. A 168, (1992) 256.
 [6] A. Nakayashiki, Int. J. Mod. Phys. A 9, (1994) 5673.
 [7] M. Jimbo and T. Miwa, J. Phys. A 29 (1996) 2923.
 [8] N. Kitanine, J. M. Maillet and V. Terras, Nucl. Phys. B 567 (2000) 554.
 [9] H.E. Boos and V.E. Korepin, J. Phys. A 34 (2001) 5311.</p> | <p>[10] H.E. Boos and V.E. Korepin, <i>Integrable models and Beyond</i>, edited by M. Kashiwara and T. Miwa (Birkhäuser, Boston, 2002); hep-th/0105144.
 [11] H.E. Boos, V.E. Korepin, Y. Nishiyama and M. Shiroishi, J. Phys. A 35 (2002) 4443.
 [12] H.E. Boos, V.E. Korepin and F.A. Smirnov, Nucl. Phys. B. 658 (2003) 417.
 [13] K. Sakai, M. Shiroishi, Y. Nishiyama and M. Takahashi, Phys. Rev. E 67 (2003) 065101(R), cond-mat/0302564.
 [14] G. Kato, M. Shiroishi, M. Takahashi, K. Sakai, J. Phys. A: Math. Gen. 36 L337-L344 (2003).</p> |
|--|---|